

Empirical Article

The role of covid-19 anxiety and intolerance of uncertainty in predicting resilienceYAGMUR BENIAN DURU,¹ VUSLAT GUNAL,² CEYDA YALCIN AGAOGLU³  and CEMRE TATLI⁴¹*Ministry of Education, School Counselor, Master Student in Hasan Kalyoncu University, Gaziantep, Turkey*²*Master Student in Hasan Kalyoncu University, Gaziantep, Turkey*³*Ministry of Education, School Counselor, PhD Student in Hasan Kalyoncu University, Kahramanmaraş, Turkey*⁴*Faculty of Education, Hasan Kalyoncu University, Gaziantep, Turkey*Duru, Y. B., Gunal, V., Agaoglu, C. Y. & Tatlı, C. (2022). The role of covid-19 anxiety and intolerance of uncertainty in predicting resilience. *Scandinavian Journal of Psychology*.

Resilience is one of the protective factors for mental health. The aim of this study is to reveal the role of COVID-19 anxiety and intolerance of uncertainty in predicting resilience in adulthood. To this end, data were collected from 533 adults (359 women-174 men). A hierarchical regression analysis was performed, controlling for the effects of gender and presence of a chronic disease on the prediction of resilience. Results showed that intolerance of uncertainty and COVID-19 anxiety were statistically significant predictors of resilience. Gender and presence of a chronic disease were not statistically significant in predicting resilience. These results indicate that intolerance of uncertainty plays an important role in explaining resilience.

Key words: COVID-19 anxiety, intolerance of uncertainty, pandemic, resilience.

Ceyda Yalcin Agaoglu, Ministry of Education, School Counselor, Kahramanmaraş, 46100, Turkey. e-mail: ceyda.ylcn@gmail.com

INTRODUCTION

COVID-19, which has spread rapidly around the world from China since the last months of 2019, has caused a global pandemic and was declared a pandemic in March 2020 (World Health Organization [WHO], 2020a). This virus causes a course that requires intensive treatment and even death (Fu, Wang, Yuan *et al.*, 2020). To protect public health and stop the spread of the virus, countries have taken measures such as isolation, quarantine, and closures (WHO, 2020b). Changes such as the high rate of spread of the disease, restrictions due to the measures taken, lifestyle changes (Haleem, Javaid & Vaishya, 2020), economic problems (Qian & Fan, 2020) have also negatively affected the mental health of individuals (Khan, Mamun, Griffiths & Ullah, 2020). During the pandemic, symptoms such as insomnia (Pappa, Ntella, Giannakas, Giannakoulis, Papoutsis & Katsaounou, 2020), depression, anxiety (Choi, Hui & Wan, 2020; Hyland, Shevlin, McBride *et al.*, 2020) and Post-traumatic stress disorder (Wang *et al.*, 2020) increased. Stress is predicted to increase the longer the pandemic lasts, leading to depression, anxiety, and uncertainty about the future (De Souza, Mohandas & Javed, 2020).

The pandemic COVID-19 brought financial and moral losses for many people. Financial, psychological, and physiological difficulties have arisen mainly due to people's worries about getting sick and losing their relatives. It has been observed that people's reactions to challenging life events such as pandemics vary widely. While some people are intimidated, others are able to accept the situation and bounce back in a short period of time. This underscores the importance of examining the factors that can protect mental health during the COVID-19 (Magson, Freeman, Rapee, Richardson, Oar & Fardouly, 2021). Indeed, it has been observed that individuals who are able to cope with the stress caused by the pandemic show fewer such symptoms

(Arkan, 2021). The ability of individuals to adapt quickly in response to such stressful life experiences is defined as resilience (Gürgan, 2006). These such traits can be learned and improved, which enable individuals to survive and overcome existing difficulties despite difficult life conditions (Gürgan, 2006; Higgins, 1994). Some individuals are able to survive and successfully continue their normal lives despite encountering a problem (Gürgan, 2006; Masten, 2001). This is an important issue with regard to psychological problems during and after the pandemic COVID-19 (Killgore, Taylor, Cloonan & Dailey, 2020).

Resilience is one of the protective factors that enable overcoming these difficulties (Naeem, Irfan & Javed, 2020; Polizzi, Lynn & Perry, 2020) and plays an important role in protecting mental health both during and after the pandemic (Killgore *et al.*, 2020). Resilience involves recovering quickly in the face of challenging events and emerging stronger from stressful and stressful events (Henderson, 2012; Jackson, Firtko & Edenborough, 2007). Resilient individuals not only adapt to difficulties, but also strive to improve by improving (Jakovljevic, 2018). During the COVID-19 pandemic, people's resilience may have decreased while their vulnerability to external factors increased (Sadati, Lankarani & Bagheri Lankarani, 2020).

There is a negative relationship between resilience and mental health (Huang, Wu, Wu, Yang, Zheng & Wu, 2020), and resilience is found to be a protective factor for mental health (Reyes, Constantino, Cross, Tan, Bombard & Acupan, 2019). On the other hand, the fact that the pandemic has been ongoing for almost two years, the recent increase in the transmission rate due to viral mutations, the lack of a definitive treatment method, the incompleteness of vaccine trials, the uncertainty of when the pandemic will end, especially the information about the epidemic and vaccines have reached the public. Factors such as the rapid spread of uncertainty are a source of fear, anxiety, and worry for

people who inherently avoid uncertainty and want to know about their future, and avoid risk. COVID-19 and the COVID-19 process triggered by the pandemic have made this uncertainty unbearable. Many people experience uncertainty about the present or the future in their lives (Sarıçam, 2014). Studies have shown that people with high intolerance of uncertainty perceive these uncertainties as fear, anxiety, and worry (Belge, 2019; Dugas, Gosselin & Ladouceur, 2001; Dugas, Hedayati, Karavidas, Buhr, Francis & Phillips, 2005; Karataş & Uzun, 2018; Kilit, Dönmezler, Erensoy & Berkol, 2020; Öz-Ceviz, Tektaş, Basmacı & Tektaş, 2020; Yıldız & Güllü, 2018). The increase in the value of intolerance of uncertainty predicts the increase in the level of anxiety (Yüksel, 2014). It is known that as intolerance of uncertainty increases, effective coping strategies for stress decrease (Coşkun, 2019). The fact that unexpected events are a risk factor for resilience (Weick & Sutcliffe, 2007) makes it important to assess resilience during the COVID-19. The uncertainty created by this unexpected process can be managed with intolerance of uncertainty. Uncertainty is a psychological state of unpredictability (Kuang & Wilson, 2017). Intolerance of uncertainty is defined as a cognitive bias that affects how a person perceives, interprets, and responds to uncertain situations, and increases anxiety (Dugas *et al.*, 2005; Fergus & Bardeen, 2013). According to Sexton, Norton, Walker and Norton (2003) intolerance of uncertainty has been shown to have an affect on anxiety and generalized fear. Also intolerance of uncertainty; it is also associated with psychological problems (Smith, Twohy & Smith, 2020), mood states such as anxiety and sadness (Meeten, Dash, Scarlet & Davey, 2012), depression, health anxiety, and maladaptive coping strategies (Rettie & Daniels, 2020). In the study of Sarıçam, Deveci and Ahmetoglu (2020), there is a negative relationship between intolerance of uncertainty and resilience. It was observed that the uncertainties in the COVID-19 have negative effects on people's mental health (Ornell, Schuch, Sordi & Kessler, 2020). In this context, it can be said that intolerance of uncertainty is a predictor of COVID-19 related worries (Voitsidis, Gliatas, Bairachtari *et al.*, 2020).

During the pandemic, individuals started to feel COVID-19 anxiety, with triggers including fear of getting sick and uncertainty (Lee, 2020). COVID-19 anxiety, coronaphobia (Asmundson & Taylor, 2020), and fear of COVID-19 (Ahorsu, Lin, Imani *et al.*, 2020), which are all associated with mental health problems (Fitzpatrick, Carson & Weisz, 2021), such as depression, anxiety, and stress (Satici, Gocet-Tekin, *et al.*, 2020). Fear of infection COVID-19 may increase the likelihood of developing mental health problems (Dsouza, Quadros, Hyderabadwala & Mamun, 2020). Management strategies and other means are very important to deal with this anxiety (Allen, Tsao, Seidman, Ehrenreich-May & Zeltzer, 2012). The previous studies stated that resilience and COVID-19 anxiety are related (Satici, Kayis, *et al.*, 2020), and resilience is considered as a protective factor (Vos, Habibović, Nyklíček, Smeets & Mertens, 2021).

It is found that as intolerance of uncertainty increases, individuals experience problems in their lives (Geçgin & Sahranc, 2017) and their anxiety levels increase (Rettie & Daniels, 2020). As a result of all these studies, it is inevitable that uncertainty has negative effects on individuals' psychological

well-being. All these findings highlight the relationship between intolerance of uncertainty and resilience.

Global pandemics have occurred throughout history. In the last 20 years in particular, we can speak of SARS and influenza pandemics as infectious diseases (Preti, Di Mattei, Perego *et al.*, 2020). In addition to research on the physical and psychological effects of infectious diseases such as SARS (Wu, Chan & Ma, 2005), 3,166 articles dealing with pandemics requiring quarantine such as SARS, Ebola, H1N1 influenza, Middle East Respiratory Syndrome (MERS), and equine influenza, it has been observed that negative psychological effects such as casualties, inadequate information and stigmatization (Brooks *et al.*, 2020). The results of previous pandemics have shown that the impact on mental health lasts longer than the pandemic itself, its prevalence is higher and calculating the psychological impact is quite difficult (Shigemura, Ursano, Morganstein, Kurosawa & Benedek, 2020). For this reason, it is important to determine the protective factors related to people's resilience during the pandemic COVID-19 in order to promote people's mental health both in the short term and after the pandemic.

As suggested in previous studies, based on the association between psychological symptoms, unexpected life events, and resilience in the COVID-19 process (Huang *et al.*, 2020), we hypothesized that both COVID-19 anxiety and intolerance of uncertainty would have predictive value for resilience. Similarly, we expected that gender and presence of a chronic disease would predict resilience, as previously suggested (Bonanno, Galea, Bucciarelli & Vlahov, 2007). Resilience was hypothesized to reduce COVID-19 anxiety and to be associated with lower intolerance of uncertainty.

METHODS

Procedure

The study has taken place during the COVID-19 pandemic. The study began after the college ethics committee approval. The scales were created in Google form and sent to participants via Whatsapp and other social media.

Participants

In order to reach the participants, the data collection tool prepared in Google form was shared on WhatsApp and Instagram social media. Snowball sampling method was used while collecting the participants. Data were collected for 4 weeks from individuals aged 18 and over who voluntarily participated in the study. Participants consisted of 673 individuals over the age of 18, 471 women and 202 men. While determining the study group, criterion sampling was used and the participants were limited according to the following criteria:

- having been diagnosed with COVID-19, Presence of a chronic disease and being hospitalized (Bolig & Weddle, 1988; Hobfoll & Lerman, 1988; Wells & Schwebel, 1987);
- being quarantined in the process of the COVID-19 pandemic;
- negative life events (Masten, Best & Gamezy, 1990; Masten, Best & Gamezy, 1999; Masten, Neemann & Andenas, 1994);
- losing one or both parents in an earlier period (Dryfoos, 1990; Rutter, 1980);
- disease (Dryfoos, 1990; Rutter, 1980); and

- having parents who live apart or divorced or living with a single parent (Grych & Fincham, 1997; Hetherington & Stanley-Hagan, 1999; Mulholland, Watt, Philpott & Sarlin, 1991). The study group took its final form as 533 individuals, 359 women and 174 men, over the age of 18. The minimum sample size for the study is “50 + 8 m” as suggested by Tabachnick and Fidell (2013). In this study, *m* (number of independent variables) is 2. In this regard, the minimum number for the study sample was found to be 66.

Demographic information: in the study, a personal information form was used and demographic information including age, gender, education level, job, marital status and having or not having children were obtained. The form, consisting of 11 items in total, also covers questions related to the inclusion criteria.

Coronavirus anxiety: Coronavirus Anxiety Scale was developed by Lee (2020) so as to measure disruptive anxiety cases and anxiety symptoms in individuals during the COVID-19. The scale, which consists of five items and a single factor is a five-point Likert type scale. In the original form of the scale, the Cronbach alpha coefficient was found to be 0.93. The adaptation studies of the scale into Turkish were performed by Akkuzu, Yumuşak, Karaman, Ladikli, Türkkân and Bahadır, (2020). In the study, the Cronbach alpha internal consistency coefficient was calculated as 0.81. Within the scope of this study, the Cronbach alpha internal consistency coefficient obtained from the Coronavirus Anxiety Scale was found to be 0.76.

Intolerance of uncertainty: the Intolerance of Uncertainty Scale (IUS-12), developed by Carleton, Norton and Asmundson (2007), is a scale enabling adults to evaluate themselves. It is a five-point Likert type scale and consists of 12 items in total. It has two sub-dimensions as Prospective Anxiety and Inhibitory Anxiety. In IUS-12, item numbered 1 is reverse coded. The highest and lowest scores that can be obtained from the scale range from 12 to 60. A high score indicates a high degree of intolerance of uncertainty. The internal consistency coefficient was found to be 0.91. The adaptation of the scale into Turkish was carried out by Sarıçam, Erguvan, Akın and Akça (2014). Internal consistency was 0.88; and it was found to be 0.84 for Prospective Anxiety and 0.77 for Inhibitory Anxiety.

Resilience: the Adult Resilience Scale was developed by Ryan and Caltabiano (2009) to measure the resilience levels of individuals in adulthood. The scale, which consists of 25 items, has five sub-dimensions. These are as follows: Factor I: Self-efficacy, Factor II: Family and Social Networks, Factor III: Perseverance, Factor IV: Internal Locus of Control, Factor V: Coping and Adaptation. It is a five-point Likert-type scale and includes reverse-coded items. After the items numbered 3, 5, 6, 10, 13, 20, 22, and 23 are reversely coded, all the items are added together to give the total resilience score. Cronbach's alpha value was found to be 0.87. The adaptation of the scale into Turkish was conducted by Savi-Çakar, Karataş and Çakır (2014). The internal consistency coefficient obtained from the adaptation study was calculated as 0.71. The Cronbach alpha coefficient for this study was found to be 0.77.

Data statistics

First, normality assumptions were tested. Group differences were tested on was performed on the variables of age and chronic disease. Hierarchical regression was performed to determine the roles of intolerance of uncertainty and COVID-19 anxiety in predicting resilience. In the analysis, first of all, the variables of gender and presence of a chronic disease were taken as control variables in the first model. In the second stage, COVID-19 anxiety and then finally the intolerance of uncertainty variable were added to the model.

With the purpose of determining the suitability of the data for hierarchical regression analysis, VIF and tolerance values were examined to determine the problem of multicollinearity between the variables. Having tolerance values above 10 and VIF values below 10 indicate that there is no multicollinearity problem (Pallant, 2011).

Considering the variables of gender and presence of a chronic disease, VIF value was found to be 1.003 in the first model and the tolerance value was 0.99 for both variables. In the second model, in which the COVID-19

anxiety was included, VIF value varied between 1.00–1.04, while the tolerance value was between 0.96 and 0.99. Finally, in the third model, which included intolerance of uncertainty, VIF value varied between 1.00–1.09 and the tolerance value varied between 0.91 and 0.99.

RESULTS

The mean, standard deviation and bivariate correlations of the variables are shown in Table 1. In order to test whether there is a difference between the means according to gender, independent groups *t*-test was performed. Gender has a statistically significant effect on COVID-19 anxiety ($t[533] = 4.433, p < 0.05$). Women ($\bar{x} = 1.90$) are more concerned about COVID-19 than men ($\bar{x} = 0.96$). There was no statistically significant effect of gender on the variables of resilience and intolerance of uncertainty. Before hierarchical regression analysis, correlations between variables were examined. There was a low negative ($r = -0.19, p < 0.05$) relationship between resilience and COVID-19 anxiety and a moderately negative ($r = -0.62, p < 0.01$) relationship between resilience and intolerance of uncertainty.

While the hierarchical multiple regression results (Table 2) do not contribute significantly to the gender regression model in the first model ($\beta = 0.03, p > 0.05$), they do make a significant contribution to the presence of a chronic disease model ($\beta = 0.09, p > 0.05$). This model explains 10% of resilience ($F_{[2,530]} = 2.63, p < 0.005$). The gender variable, which had no effect in the first model, did not make a significant contribution in the second model either ($\beta = 0.07, p > 0.05$). The COVID-19 anxiety included in the second model is a significant ($\beta = -0.20, p > 0.001$) predictor ($F_{[3,529]} = 9.59, p < 0.005$), explaining 5% of the change in resilience.

Presence of a chronic disease, which was significant in the first two models, contributed significantly to the third model ($\beta = 0.06, p > 0.05$), while the gender variable did not contribute significantly to this model ($\beta = 0.02, p > 0.05$). The intolerance of uncertainty variable added in the last model is a significant predictor of the model ($\beta = -0.60, p > 0.001$) and explains 39% of the change in resilience ($F_{[4,528]} = 85.52, p < 0.005$). In the light of these findings, it can be said that intolerance of uncertainty has a full mediator effect between COVID-19 anxiety and resilience.

TABLE 1. Means, standard deviations and correlations between the variables of the study

	1.	2.	3.	4.	Mean	SD
1. Resilience	1.00	−0.62**	−0.19*	0.09*	67.79	10.02
2. Intolerance of uncertainty		1.00	0.21**	−0.05	40.40	9.21
3. COVID-19 anxiety			1.00	0.00	1.60	2.34
4. Presence of a chronic disease				1.00	1.89	0.30

* $p < 0.05$,

** $p < 0.01$.

TABLE 2. Results of hierarchical regression analyses regressing resilience

Predictors	Model 1	Model 2	Model 3
Presence of a chronic disease	0.09	0.09	0.06
Gender	0.03	0.07	0.02
COVID-19 anxiety		-0.20*	-0.07
Intolerance of uncertainty			-0.60*
ΔR^2	0.01	0.05	0.39*
Total adjusted R ²	0.00	0.04	0.38*

* $p < 0.01$.

DISCUSSION

Considering the results of the study, intolerance of uncertainty and fear of COVID-19 negatively predict resilience. Resilience affects COVID-19 anxiety both directly and through intolerance of uncertainty. Based on this result, a person with high resilience can be expected to have less COVID-19 anxiety. This result is consistent with previous findings (e.g., Kasapoğlu, 2020). Another finding of the study is that resilience negatively predicts intolerance of uncertainty and COVID-19 anxiety. It was found that intolerance of uncertainty has a higher predictive value. One of the most important findings of this study is that intolerance of uncertainty fully mediates the relationship between resilience and COVID-19 anxiety. Previous studies showed that intolerance of uncertainty, fear of COVID-19 and mediation studies with different variables. Intolerance of uncertainty was found to play a mediating role in the relationship between fear of COVID-19 and positivity (Bakioğlu, Korkmaz & Ercan, 2020). Deniz (2021) examined the mediating role of intolerance of uncertainty and fear of COVID-19 in the relationship between self-concept and well-being. It has been shown that individuals with higher self-concept related to intolerance of uncertainty have lower fear of COVID-19.

There is some evidence showing the relationship between resilience and intolerance of uncertainty (DiTrani, Mariani, Ferri, De Berardinis & Frigo, 2021) and the relationship between resilience, and COVID-19 anxiety (Labrague & De Los Santos, 2020). There is also a study in which COVID-19 anxiety predicts resilience (Karataş & Tagay, 2021). Based on the negative relationship between resilience and intolerance of uncertainty in the pre-pandemic period (Lee, 2019; Mitmansgruber, Smrekar, Rabanser, Beck, Eder & Ellemunter, 2016), it can be said that this relationship is independent of the COVID-19. Given the protective role of resilience to negative and unexpected life events such as COVID-19 (Naem *et al.*, 2020) and given the role of COVID-19 anxiety, studies of resilience, and intolerance of uncertainty should be conducted to support the mental health of individuals' post-pandemic.

This study examined the predictive role of intolerance of uncertainty and COVID-19 anxiety in predicting adult resilience levels during the COVID-19. In addition, resilience, intolerance of uncertainty and COVID-19 anxiety were assessed by gender and the presence of a chronic disease. Women were found to experience more COVID-19 anxiety.

In the study, COVID-19 anxiety was found to be more prevalent in women. Considering similar findings in different

cultures (Hyland *et al.*, 2020; Lee, 2020; Miguel-Puga, Cooper-Bribiesca, Avelar-Garnica *et al.*, 2020; Öz-Ceviz *et al.*, 2020; Solomou & Constantinidou, 2020), it is reasonable to assume that this is independent of culture. Furthermore, this situation is thought to be related to women being more vulnerable to stress-related disorders such as post-traumatic stress disorder and anxiety disorders (Li & Graham, 2017). In the literature, there are studies showing that men have higher levels of resilience (Bonanno *et al.*, 2007; Masood, Masud & Mazahir, 2016) as well as studies indicating that there is no difference between genders (Isaacs, Mota, Tsai *et al.*, 2017; Karataş & Tagay, 2021; Ünver & Güloğlu, 2021). In general, women are physically weaker, have lower immunity and are potentially more likely to get sick, but they adapt better to environmental conditions and live longer (Overfield, 2018). Gender roles are another factor that can be discussed in this finding. Because women are perceived in society as weaker, in need of protection and fragile, they are in a culturally acceptable position to seek help and treatment. Since men need to look stronger, be brave and take on the role of protector, the duration of disease may be longer as they experience a delayed process while seeking help (Bakioğlu *et al.*, 2020). Resilience is thought to be related to difficult life events rather than gender. Resilience can be considered as a force that occurs after traumatic life events such as death, chronic disease, divorce, natural disasters or epidemic diseases, rather than a situation that occurs or is triggered during daily life events (Yazıcı-Çelebi, 2020).

The COVID-19 pandemic is the most important health problem in our country and in the whole world. In particular, people over 60 years of age, cancer, diabetes, heart, hypertension, chronic liver, and kidney diseases, obesity, Parkinson's disease, MS patients, chronic lung diseases, bronchitis or asthma, people receiving immuno suppressive therapy, and people with immunodeficiency are among the risk groups (Sofulu, Uran, Avdal & Tokem, 2020). Examining the literature; studies on resilience and chronic disease, resilience and depression, anxiety, somatization, heart disease, etc., highlight the negative relationship between mental and physical disease (Cal, De Sá, Glustak & Santiago, 2015; Toukhsati, Jovanovic, Dehghani, Tran, Tran & Hare, 2017). On the contrary, even in vulnerable populations such as adults with chronic diseases, it is observed that individuals can show resilience or increase their existing level of resilience in pandemic conditions such as COVID-19 (Davis, McElroy-Heltzel, Lemke *et al.*, 2021).

On the other hand, there is no difference between genders in the variables of resilience and intolerance of uncertainty. The literature mentions gender differences in resilience, intolerance of uncertainty and COVID-19 anxiety. However, in this study, the presence of a chronic disease did not significantly affect resilience. The reason why there is no relationship between presence of a chronic disease and resilience in this study, contrary to the literature, is that the selected sample was not resilience prone and the group that was not resilience prone was eliminated at the beginning of the study and a homogeneous group was studied with a homogeneous group that experienced negative life events (COVID-19 disease, being hospitalized, presence of a chronic disease, etc.) could be. However, as resilience develops over time and events, the individual's past experiences may be

effective in increasing or decreasing resilience in future life stages (Gheshlagh, Ebadi, Dalvandi, Rezaei & Tabrizi, 2017). In this regard, the increase in age of the individuals participating in the study leads to the belief that the relationship between presence of a chronic disease and resilience is also increasing.

Limitations and future directions

This study has some limitations. Although COVID-19 has been an ongoing process for more than a year, the data were collected within the first six months of the COVID-19 period. Due to the fact that the data were collected online, rural areas without internet access could not be reached. This data has limited generalizability because it was only collected from individuals who have a technology background and use the Internet. Furthermore, our study is a cross-sectional study. This represents one of the limitation of the study, because it is predicted that the pandemic COVID-19 will be a long and dynamic process and that both physical and psychological effects will occur over a long period of time. For this reason, longitudinal studies are needed to investigate the problem thoroughly. The fact that resilience is a protective factor suggests that related variables should be studied in depth, so future studies that incorporate qualitative data in groups of different ages and socioeconomic levels will contribute to the literature.

This study was conducted on adults over 18 years of age. By conducting research with school-aged individuals affected by the pandemic, it is possible to examine how intolerance of uncertainty and COVID-19 anxiety predict the level of resilience of these individuals. This study examined individuals between the ages of 18 and 65 years, indicating that there was no homogeneous group to be obtained in terms of life task characteristics. Although the age of 18 is legally recognized as an adult, it is believed that resilience in particular makes a difference between the experiences of an 18-year-old and a 65-year-old. In this regard, it is predicted that work will have a more homogeneous structure in periods suitable for developmental tasks. The variables intolerance of uncertainty and COVID-19 anxiety can be considered in efforts to increase or protect the resilience of individuals during and after the pandemic.

Studying with a sample in which the criteria for resilience related to negative life events were determined only in the context of COVID-19 may be beneficial for specific results. At the same time, the variable age has been reported in the literature as both a protective and a risk factor.

This study discusses variables that are expected to negatively affect resilience. Considering the many negative emotional and behavioral consequences of COVID-19 anxiety and intolerance of uncertainty for individuals, preventive studies should be conducted to protect the spiritual and mental health of society during the pandemics. To this end, both studies that prevent negative effects and studies that support individuals' strengths should be also taken into consideration.

CONCLUSIONS

Intolerance of uncertainty has a full mediator effect between COVID-19 anxiety and resilience. While there is no gender

difference in the resilience and intolerance of uncertainty variables, women seem to experience higher COVID-19 anxiety.

As a highly contagious disease, COVID-19 was recognized as a pandemic worldwide (WHO, 2020a). Delays in diagnosing and treating the disease and people's anxiety about returning to their daily lives have created an environment of uncertainty. In short, the increase in COVID-19 anxiety leads to an increase in intolerance of uncertainty (Kasapoğlu, 2020). This is because it is not easy to regulate emotions when there is intolerance of uncertainty (Deniz, 2021). Resilience, on the other hand, is considered a protective factor for all these experiences and ensures that individuals with resilience are more positive and tolerant of uncertainty COVID-19.

There is no difference between genders in the variables of resilience and intolerance of uncertainty. Women experience higher COVID-19 anxiety. Intolerance of uncertainty and COVID-19 anxiety are predictive factors for resilience. Intolerance of uncertainty has a full mediator effect between COVID-19 anxiety and resilience.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are openly available in [repository name e.g. "figshare"] at [http://doi.org/\[doi\], reference number \[reference number\]](http://doi.org/[doi], reference number [reference number]).

REFERENCES

- Ahorsu, D.K., Lin, C.Y., Imani, V., Saffari, M., Griffiths, M.D. & Pakpour, A.H. (2020). The fear of COVID-19 scale: Development and initial validation. *International Journal of Mental Health and Addiction*, 1–9. <https://doi.org/10.1007/s11469-020-00270-8>.
- Allen, L. B., Tsao, J. C., Seidman, L. C., Ehrenreich-May, J. & Zeltzer, L. K. (2012). A unified, transdiagnostic treatment for adolescents with chronic pain and comorbid anxiety and depression. *Cognitive and Behavioral Practice*, 19, 56–67.
- Akkuzu, H., Yumuşak, F.N., Karaman, G., Ladikli, N., Türkkan, Z. & Bahadır, E. (2020). Koronavirüs Kaygı Ölçeği'nin Türkçe güvenilirlik ve geçerlik çalışması. *Kıbrıs Türk Psikiyatri ve Psikoloji Dergisi*, 2, 63–67.
- Arıkan, G. (2021). COVID-19 salgınında stres: Olumsuz, olumlu sonuçları ve önleyici müdahalenin olası rolü. *Psikiyatride Güncel Yaklaşımlar*, 13, 135–145.
- Asmundson, G. J. & Taylor, S. (2020). Coronaphobia revisited: A state-of-the-art on pandemic-related fear, anxiety, and stress. *Journal of Anxiety Disorders*, 76, <https://doi.org/10.1016/j.janxdis.2020.102326>
- Bakioğlu, F., Korkmaz, O. & Ercan, H. (2020). Fear of COVID-19 and positivity: Mediating role of intolerance of uncertainty, depression, anxiety, and stress. *International Journal of Mental Health and Addiction*, 19, 2369–2382.
- Belge, J. (2019). *Bir grup yetişkinde depresif semptomlar, anksiyete semptomları ve belirsizliğe tahammülsüzlük arasındaki ilişkinin belirlenmesi [Yayınlanmamış Yüksek Lisans Tezi]*. İstanbul: İstanbul Gelişim Üniversitesi.
- Bolig, R. & Weddle, K.D. (1988). Resiliency and hospitalization of children. *Children's Health Care*, 16, 255–260.
- Bonanno, G.A., Galea, S., Bucchiarelli, A. & Vlahov, D. (2007). What predicts psychological resilience after disaster? The role of demographics, resources, and life stress. *Journal of Consulting and Clinical Psychology*, 75, 671–682.
- Brooks, S.K., Webster, R.K., Smith, L.E., Woodland, L., Wessely, S., Greenberg, N. et al. (2020). The psychological impact of quarantine and how to reduce it: Rapid review of the evidence. *The Lancet*, 395, 912–920.

- Cal, S.F., De Sá, L.R., Glustak, M.E. & Santiago, M.B. (2015). Resilience in chronic diseases: A systematic review. *Cogent Psychology*, 2, 1–9.
- Carleton, R.N., Norton, M.A. & Asmundson, G.J.G. (2007). Fearing the unknown: A short version of the intolerance of uncertainty scale. *Journal of Anxiety Disorders*, 21, 105–117.
- Choi, E.P.H., Hui, B.P.H. & Wan, E.Y.F. (2020). Depression and anxiety in Hong Kong during COVID-19. *International Journal of Environmental Research and Public Health*, 17, 3740. <https://doi.org/10.3390/ijerph17103740>.
- Coşkun, E. (2019). *Duygusal zeka ve belirsizliğe tahammülsüzlüğün stresle başa çıkma tarzlarına etkisinin incelenmesi [Yayınlanmamış Yüksek Lisans Tezi]* (pp. 1–144). İstanbul: Fatih Sultan Mehmet Vakıf Üniversitesi.
- Davis, E.B., McElroy-Heltzel, S.E., Lemke, A.W., Cowden, R.G., VanderWeele, T.J., Worthington, E.L., Jr. et al. (2021). Psychological and spiritual outcomes during the COVID-19 pandemic: A prospective longitudinal study of adults with chronic disease. *Health Psychology*, 40, 347–356.
- De Souza, A., Mohandas, E. & Javed, A. (2020). Psychological interventions during COVID-19: Challenges for low and middle income countries. *Asian Journal of Psychiatry*, 51, 102128. <https://doi.org/10.1016/j.ajp.2020.102128>.
- Deniz, M.E. (2021). Self-compassion, intolerance of uncertainty, fear of COVID-19, and well-being: A serial mediation investigation. *Personality and Individual Differences*, 177, 110824. <https://doi.org/10.1016/j.paid.2021.110824>.
- DiTrani, M., Mariani, R., Ferri, R., De Berardinis, D. & Frigo, M.G. (2021). From resilience to burnout in healthcare workers during the COVID-19 emergency: The role of the ability to tolerate uncertainty. *Frontiers in Psychology*, 12, 987. <https://doi.org/10.3389/fpsyg.2021.646435>.
- Dryfoos, J.G. (1990). *Adolescents at risk: Prevalence and prevention*. Oxford: Oxford University Press.
- Dsouza, D.D., Quadros, S., Hyderabadwala, Z.J. & Mamun, M.A. (2020). Aggregated COVID-19 suicide incidences in India: Fear of COVID-19 infection is the prominent causative factor. *Psychiatry Research*, 290, 113145. <https://doi.org/10.1016/j.psychres.2020.113145>.
- Dugas, M.J., Gosselin, P. & Ladouceur, R. (2001). Intolerance of uncertainty and worry: Investigating specificity in a nonclinical sample. *Cognitive Therapy and Research*, 25, 551–558.
- Dugas, M.J., Hedayati, M., Karavidas, A., Buhr, K., Francis, K. & Phillips, N.A. (2005). Intolerance of uncertainty and information processing: Evidence of biased recall and interpretations. *Cognitive Therapy and Research*, 29, 57–70.
- Fergus, T.A. & Bardeen, J.R. (2013). Anxiety sensitivity and intolerance of uncertainty: Evidence of incremental specificity in relation to health anxiety. *Personality and Individual Differences*, 55, 640–644.
- Fitzpatrick, O., Carson, A. & Weisz, J.R. (2021). Using mixed methods to identify the primary mental health problems and needs of children, adolescents, and their caregivers during the coronavirus (COVID-19) pandemic. *Child Psychiatry & Human Development*, 52, 1082–1093.
- Fu, L., Wang, B., Yuan, T., Chen, X., Ao, Y., Fitzpatrick, T. et al. (2020). Clinical characteristics of coronavirus disease 2019 (COVID-19) in China: a systematic review and meta-analysis. *Journal of Infectious Diseases*, 80, 656–665.
- Geçgin, F.M. & Sahraç, Ü. (2017). Belirsizliğe tahammülsüzlük ve psikolojik iyi oluş arasındaki ilişki. *Sakarya University Journal of Education*, 7, 739–755.
- Gheshlagh, R.G., Ebadi, A., Dalvandi, A., Rezaei, M. & Tabrizi, K.N. (2017). A systematic study of resilience in patients with chronic physical diseases. *Nursing and Midwifery Studies*, 6, e36401. <https://doi.org/10.5812/nmsjournal.36401>.
- Grych, J.H. & Fincham, F.D. (1997). Children's adaptation to divorce. In S. A. Wolchik & I. N. Sandler (Eds.), *Handbook of children's coping*. Issues in Clinical Child Psychology (pp. 159–193). Boston, MA: Springer.
- Gürkan, U. (2006). Yılmazlık ölçeği (yö): Ölçek geliştirme, güvenilirlik ve geçerlik çalışması. *Ankara Üniversitesi Eğitim Bilimleri Fakültesi Dergisi*, 39, 45–74.
- Haleem, A., Javaid, M. & Vaishya, R. (2020). Effects of COVID 19 pandemic in daily life. *Current Medicine Research and Practice*, 10, 78–79.
- Henderson, N. (2012). Resilience in schools and curriculum design. In M. Ungar (Ed.), *The social ecology of resilience* (pp. 297–306). New York: Springer.
- Hetherington, E.M. & Stanley-Hagan, M. (1999). The adjustment of children with divorced parents: A risk and resiliency perspective. *Journal of Child Psychology and Psychiatry*, 40, 129–140.
- Higgins, G.O. (1994). *Resilient adults: Overcoming a cruel past*. San Francisco: Jossey-Bass.
- Hobfoll, S.E. & Lerman, M. (1988). Personal relationships, personal attributes, and stress resistance: Mothers' reactions to their child's illness. *American Journal of Community Psychology*, 16, 565–589.
- Huang, Y., Wu, R., Wu, J., Yang, Q., Zheng, S. & Wu, K. (2020). Psychological resilience, self-acceptance, perceived social support and their associations with mental health of incarcerated offenders in China. *Asian Journal of Psychiatry*, 52, 102166. <https://doi.org/10.1016/j.ajp.2020.102166>.
- Hyland, P., Shevlin, M., McBride, O., Murphy, J., Karatzias, T., Bentall, R.P. et al. (2020). Anxiety and depression in the Republic of Ireland during the COVID-19 pandemic. *Acta Psychiatrica Scandinavica*, 142, 249–256.
- Isaacs, K., Mota, N.P., Tsai, J., Harpaz-Rotem, I., Cook, J.M., Kirwin, P.D. et al. (2017). Psychological resilience in US military veterans: A 2-year, nationally representative prospective cohort study. *Journal of Psychiatric Research*, 84, 301–309.
- Jackson, D., Firtko, A. & Edenborough, M. (2007). Personal resilience as a strategy for surviving and thriving in the face of workplace adversity: A literature review. *Journal of Advanced Nursing*, 60, 1–9.
- Jakovljevic, M. (2018). Empathy, sense of coherence and resilience: Bridging personal, public and global mental health and conceptual synthesis. *Psychiatria Danubina*, 30, 380–384.
- Karataş, Z. & Tagay, Ö. (2021). The relationships between resilience of the adults affected by the covid pandemic in Turkey and COVID-19 fear, meaning in life, life satisfaction, intolerance of uncertainty and hope. *Personality and Individual Differences*, 172, 110592. <https://doi.org/10.1016/j.paid.2020.110592>.
- Karataş, Z. & Uzun, K. (2018). Belirsizliğe tahammülsüzlüğün yordayıcısı olarak endişe ile ilgili olumlu ve olumsuz inançlar. *Kastamonu Education Journal*, 26, 1267–1276.
- Kasapoğlu, F. (2020). COVID-19 salgını sürecinde kaygı ile maneviyat, psikolojik sağlamlık ve belirsizliğe tahammülsüzlük arasındaki ilişkilerin incelenmesi. *Turkish Studies*, 15, 599–614.
- Khan, K.S., Mamun, M.A., Griffiths, M.D. & Ullah, I. (2020). The mental health impact of the COVID-19 pandemic across different cohorts. *International Journal of Mental Health and Addiction*, 20, 1–7.
- Kilit, Z., Dönmezler, S., Erensoy, H. & Berkol, T.D. (2020). Üniversite öğrencilerinde belirsizliğe tahammülsüzlük, endişe ve bilişsel sınav kaygısı ilişkisi. *Ortadoğu Tıp Dergisi*, 12, 262–268.
- Killgore, W.D., Taylor, E.C., Cloonan, S.A. & Dailey, N.S. (2020). Psychological resilience during the COVID-19 lockdown. *Psychiatry Research*, 291, 113216. <https://doi.org/10.1016/j.psychres.2020.113216>.
- Kuang, K. & Wilson, S.R. (2017). A meta-analysis of uncertainty and information management in illness contexts. *Journal of Communication*, 67, 378–401.
- Labrague, L.J. & De Los Santos, J.A.A. (2020). COVID-19 anxiety among front-line nurses: Predictive role of organisational support, personal resilience and social support. *Journal of Nursing Management*, 28, 1653–1661.
- Lee, J.S. (2019). Effect of resilience on intolerance of uncertainty in nursing university students. *Nursing Forum*, 54, 53–59.
- Lee, S.A. (2020). Coronavirus anxiety scale: A brief mental health screener for COVID-19 related anxiety. *Death Studies*, 44, 393–401.
- Li, S.H. & Graham, B.M. (2017). Why are women so vulnerable to anxiety, trauma-related and stress-related disorders? The potential role of sex hormones. *The Lancet Psychiatry*, 4, 73–82.
- Magson, N.R., Freeman, J.Y., Rapee, R.M., Richardson, C.E., Oar, E.L. & Fardouly, J. (2021). Risk and protective factors for prospective

- changes in adolescent mental health during the COVID-19 pandemic. *Journal of Youth and Adolescence*, 50, 44–57.
- Masood, A., Masud, Y. & Mazahir, S. (2016). Gender differences in resilience and psychological distress of patients with burns. *Burns*, 42, 300–306.
- Masten, A.S. (2001). Ordinary magic: resilience processes in development. *American Psychologist*, 56, 227–238.
- Masten, A.S., Best, K.M. & Garmezy, N. (1990). Resilience and development: Contributions from the study of children who overcome adversity. *Development and Psychopathology*, 2, 425–444.
- Masten, A.S., Hubbard, J.J., Gest, S.D., Tellegen, A., Garmezy, N. & Ramirez, M. (1999). Competence in the context of adversity: Pathways to resilience and maladaptation from childhood to late adolescence. *Development and Psychopathology*, 11, 143–169.
- Masten, A.S., Neemann, J. & Andenas, S. (1994). Life events and adjustment in adolescents: The significance of event independence, desirability, and chronicity. *Journal of Research on Adolescence*, 4, 71–97.
- Meeten, F., Dash, S. R., Scarlet, A.L.S. & Davey, G.C.L. (2012). Investigating the effect of intolerance of uncertainty on catastrophic worrying and mood. *Behaviour Research and Therapy*, 50, 690–698.
- Miguel-Puga, J.A., Cooper-Bribiesca, D., Avelar-Garnica, F.J., Sanchez-Hurtado, L.A., Colin-Martínez, T., Espinosa-Poblano, E. *et al.* (2020). Burnout, depersonalization and anxiety contribute to post-traumatic stress in frontline health workers at COVID-19 patient care, a follow up study. *Brain and Behavior*, 11, 1–9.
- Mitmansgruber, H., Smrekar, U., Rabanser, B., Beck, T., Eder, J. & Ellemunter, H. (2016). Psychological resilience and intolerance of uncertainty in coping with cystic fibrosis. *Journal of Cystic Fibrosis*, 15, 689–695.
- Mulholland, D.J., Watt, N.F., Philpott, A. & Sarlin, N. (1991). Academic performance in children of divorce: Psychological resilience and vulnerability. *Psychiatry*, 54, 268–280.
- Naem, F., Irfan, M. & Javed, A. (2020). Coping with COVID-19: Urgent need for building resilience through cognitive behaviour therapy. *Khyber Medical University Journal*, 12, 1–3.
- Ornell, F., Schuch, J.B., Sordi, A.O. & Kessler, F.H.P. (2020). Pandemic fear and COVID-19: Mental health burden and strategies. *Brazilian Journal of Psychiatry*, 42, 232–235.
- Overfield, T. (2018). *Biological variation in health and illness: Race, age, and sex*. Boca Raton: CRC Press.
- Öz-Ceviz, N., Tektaş, N., Basmacı, G. & Tektaş, M. (2020). Covid 19 pandemi sürecinde üniversite öğrencilerinin kaygı düzeylerini etkileyen değişkenlerin analizi. *Uluslararası Eğitim Araştırmacıları Dergisi*, 3, 312–329.
- Pallant, J. (2011). *SPSS survival manual*. Maidenhead, Berkshire: McGraw-Hill Education.
- Pappa, S., Ntella, V., Giannakas, T., Giannakoulis, V.G., Papoutsis, E. & Katsaounou, P. (2020). Prevalence of depression, anxiety, and insomnia among healthcare workers during the COVID-19 pandemic: A systematic review and meta-analysis. *Brain, Behavior, and Immunity*, 88, 901–907.
- Polizzi, C., Lynn, S. J. & Perry, A. (2020). Stress and coping in the time of COVID-19: Pathways to resilience and recovery. *Clinical Neuropsychiatry*, 17, 59. <https://doi.org/10.36131/CN20200204>
- Preti, E., Di Mattei, V., Perego, G., Ferrari, F., Mazzetti, M., Taranto, P. *et al.* (2020). The psychological impact of epidemic and pandemic outbreaks on healthcare workers: Rapid review of the evidence. *Current Psychiatry Reports*, 22, 1–22.
- Qian, Y. & Fan, W. (2020). Who loses income during the COVID-19 outbreak? Evidence from China. *Research in Social Stratification and Mobility*, 68, 100522. <https://doi.org/10.1016/j.rssm.2020.100522>.
- Rettie, H. & Daniels, J. (2020). Coping and tolerance of uncertainty: Predictors and mediators of mental health during the COVID-19 pandemic. *American Psychologist*, 76, 427–437.
- Reyes, A.T., Constantino, R.E., Cross, C.L., Tan, R.A., Bombard, J.N. & Acupan, A.R. (2019). Resilience and psychological trauma among Filipino American women. *Archives of Psychiatric Nursing*, 33, 177–185.
- Rutter, M. (1980). Psychosexual development. In M. Rutter (Ed.), *Scientific Foundations of Developmental Psychiatry*. London: Heinemann Medical.
- Ryan, L. & Caltabiano, M. (2009). Development of a new resilience scale: The resilience in midlife scale. *Asian Social Science*, 5, 39–51.
- Sadati, A.K.B., Lankarani, M.H. & Bagheri Lankarani, K. (2020). Risk society, global vulnerability and fragile resilience; sociological view on the corona virus outbreak. *Advances in Shiraz E-Medical Journal*, 21. <https://doi.org/10.5812/semj.102263>
- Sarıçam, H. (2014). Belirsizliğe tahammülsüzlüğün mutluluğa etkisi. *Sosyal Bilimler Dergisi*, 4, 1–12.
- Sarıçam, H., Devenci, M. & Ahmetoglu, E. (2020). The examination of hope, intolerance of uncertainty and resilience levels in parents having disabled children. *Global Journal of Psychology Research: New Trends and Issues*, 10, 118–131.
- Sarıçam, H., Erguvan, F.M., Akın, A. & Akça, M.Ş. (2014). Belirsizliğe tahammülsüzlük ölçeği (btö-12) Türkçe formu: Geçerlilik ve güvenilirlik çalışması. *Route Educational and Social Science Journal*, 1, 148–157.
- Satici, B., Gocet-Tekin, E., Deniz, M.E. & Satici, S.A. (2020). Adaptation of the fear of COVID-19 scale: Its association with psychological distress and life satisfaction in Turkey. *International Journal of Mental Health and Addiction*, 19, 1980–1988.
- Satici, S.A., Kayis, A.R., Satici, B., Griffiths, M.D. & Can, G. (2020). Resilience, hope, and subjective happiness among the turkish population: Fear of COVID-19 as a mediator. *International Journal of Mental Health and Addiction*, 1–16. <https://doi.org/10.1007/s11469-020-00443-5>.
- Savi-Çakar, F., Karataş, Z. & Çakır, M.A. (2014). Yetişkin yılmazlık ölçeği: Türk kültürüne uyarlanması. *Mehmet Akif Ersoy Üniversitesi Eğitim Fakültesi Dergisi*, 32, 22–39.
- Sexton, K.A., Norton, P.J., Walker, J.R. & Norton, G.R. (2003). Hierarchical model of generalized and specific vulnerabilities in anxiety. *Cognitive Behaviour Therapy*, 32, 82–94.
- Shigemura, J., Ursano, R.J., Morganstein, J.C., Kurosawa, M. & Benedek, D.M. (2020). Public responses to the novel 2019 coronavirus (2019-nCoV) in Japan: Mental health consequences and target populations. *Psychiatry and Clinical Neurosciences*, 74, 281. <https://doi.org/10.1111/pcn.12988>.
- Smith, B.M., Twohy, A.J. & Smith, G.S. (2020). Psychological inflexibility and intolerance of uncertainty moderate the relationship between social isolation and mental health outcomes during COVID-19. *Journal of Contextual Behavioral Science*, 18, 162–174.
- Sofulu, F., Uran, B.Ö., Avdal, E.Ü. & Tokem, Y. (2020). COVID-19 salgınında kronik hastalıklarda hemşirelik yönetimi. *İzmir Katip Çelebi Üniversitesi Sağlık Bilimleri Fakültesi Dergisi*, 5, 147–151.
- Solomou, I. & Constantinidou, F. (2020). Prevalence and predictors of anxiety and depression symptoms during the COVID-19 pandemic and compliance with precautionary measures: age and sex matter. *International Journal of Environmental Research and Public Health*, 17, 4924. <https://doi.org/10.3390/ijerph17144924>.
- Tabachnick, B.G. & Fidell, L.S. (2013). *Using multivariate statistics: International edition*. Pearson 2012. Boston: Allyn and Bacon.
- Toukhsati, S.R., Jovanovic, A., Dehghani, S., Tran, T., Tran, A. & Hare, D.L. (2017). Low psychological resilience is associated with depression in patients with cardiovascular disease. *European Journal of Cardiovascular Nursing*, 16, 64–69.
- Ünver, D. & Güloğlu, B. (2021). COVID-19 pandemisinde evli bireylerde depresyon, anksiyete ve stresin ilişkisel yılmazlık bağlamında incelenmesi. *ASEAD*, 8, 517–531.
- Voitsidis, P., Gliatas, I., Bairachari, V., Papadopoulou, K., Papageorgiou, G., Parlapani, E. *et al.* (2020). Insomnia during the COVID-19 pandemic in a Greek population. *Psychiatry Research*, 289, 113076. <https://doi.org/10.1016/j.psychres.2020.113076>.
- Vos, L.M., Habibović, M., Nyklíček, I., Smeets, T. & Mertens, G. (2021). Optimism, mindfulness, and resilience as potential protective factors for the mental health consequences of fear of the coronavirus. *Psychiatry Research*, 300, 113927. <https://doi.org/10.1016/j.psychres.2021.113927>.

- Wang, Y.X., Guo, H. T., Du, X.W., Song, W., Lu, C. & Hao, W.N. (2020). Factors associated with post-traumatic stress disorder of nurses exposed to corona virus disease 2019 in China. *Medicine*, 99. <https://doi.org/10.1097/MD.00000000000020965>
- Weick, K.E. & Sutcliffe, K.M. (2007). *Managing the unexpected: resilient performance in an age of uncertainty*. San Francisco: Jossey-Bass.
- Wells, R.D. & Schwebel, A.I. (1987). Chronically ill children and their mothers: Predictors of resilience and vulnerability to hospitalization and surgical stress. *Journal of Developmental and Behavioral Pediatrics*, 8, 83–89.
- World Health Organization. (2020a). Novel Coronavirus (2019-nCoV) Situation Report-1. Retrieved 20 September 2020 from https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200121-sitrep-1-2019-ncov.pdf?sfvrsn=20a99c10_4
- World Health Organization. (2020b). WHO Director-General's remarks at the media briefing on 2019-nCoV Retrieved 20 September 2020 from <https://www.who.int/dg/speeches/detail/who-directorgeneral-s-remarksat-the-media-briefing-on-2019-ncov-on-11-february-2020>
- Wu, K.K., Chan, S.K. & Ma, T.M. (2005). Posttraumatic stress, anxiety, and depression in survivors of severe acute respiratory syndrome (SARS). *Journal of Traumatic Stress: Official Publication of The International Society for Traumatic Stress Studies*, 18, 39–42.
- Yazıcı-Çelebi, G. (2020). COVID-19 salgınına ilişkin tepkilerin psikolojik sağlamlık açısından incelenmesi. *IBAD Journal of Social Sciences*, 8, 471–483.
- Yıldız, B. & Güllü, A. (2018). Belirsizliğe tahammülsüzlük ile aleksitimi arasındaki ilişki ve bazı sosyodemografik değişkenlere göre incelenmesi. *Eğitim Bilimleri Dergisi*, 9, 113–131.
- Yüksel, B. (2014). *Kayı Belirtilerini Açıklamada Bağlanma, Pozitif ve Negatif Duygu Düzenleme ve Belirsizliğe Tahammülsüzlük Arasındaki İlişkiyi Bütünleyici Model Arayışı [Yayınlanmamış Yüksek Lisans Tezi]*. Ankara: Hacettepe Üniversitesi.

Received 22 August 2021, Revised 11 March 2022, accepted 20 March 2022